

Separation of integer points by a hyperplane under some weak notions of discrete convexity

Takuya Kashimura*, Yasuhide Numata*[†] and Akimichi Takemura*[†]

February 2010

Abstract

We give some sufficient conditions of separation of two sets of integer points by a hyperplane. Our conditions are related to the notion of convexity of sets of integer points and are weaker than existing notions.

1 Introduction

In this paper, for a given nonempty set S of integer points in $\mathbb{R}^d$, we investigate conditions on separation of two sets $A \subset S$ and $B = S \setminus A$ by an affine hyperplane. Consideration of hyperplane separation naturally leads to various notions of discrete convexity of A and B . The hyperplane separation theorem for two disjoint convex sets in $\mathbb{R}^d$, or the Hahn-Banach separation theorem for infinite dimensional spaces, is a fundamental fact and is a logical basis for various fields such as optimization or game theory. However, because of the discreteness, separation of sets of integer points is a subtle problem. Murota ([6, 7, 8]) and his collaborators have developed the whole new field of “discrete convex analysis” and the hyperplane separation is an important motivation for the field.

*Department of Mathematical Informatics, Graduate School of Information Science and Technology, University of Tokyo.

[†]JST, CREST

In Murota's works and works by earlier authors, hyperplane separation results have been proved by imposing some nice conditions of discrete convexity of A and B . However for some problems we want to know whether separation holds under weaker conditions on S , A and B . In [2] we needed a separation result for $\mathbb{R}^2$ under "parallelogram condition" given in Definition 2.9 below. A similar result for higher dimension was conjectured in [2] and the conjecture is the motivation for the present study. As shown in Section 4 we found that there is a large gap between $\mathbb{R}^2$ and $\mathbb{R}^d$ for $d \geq 3$.

Non-convexity of a set A of integer points is usually characterized by existence of a "hole" in A . Recently active research is conducted on various notions of holes. Fano polytopes, which are motivated from algebraic geometry, are actively investigated from combinatorial viewpoint [4, 5]. They are rich sources of polytopes with a single hole in its interior. Empty lattice simplices, whose only integer points are its vertices, have been studied in many contexts (see [10] and references therein). In the field of commutative algebra and its application to algebraic statistics, holes in a semigroup generated by a set of integer points are of great interest (e.g. [3, 9]).

The organization of this paper is as follows. In Section 2 we give definitions and some preliminary facts. In Section 3 we prove that hyperplane separation holds under very weak condition for the case $S = \mathbb{Z}^d$. In Section 4 we consider finite A and B and prove separation results when $S \subset \mathbb{Z}^2$ is integrally convex and when $S \subset \mathbb{Z}^2$ is hole free. For $\mathbb{Z}^d$, $d \geq 3$, we give some counterexamples. We end the paper with discussion of some open problems in Section 5.

2 Definitions and some preliminary facts

For a given nonempty set S of integer points in $\mathbb{R}^d$, we consider some conditions which concern separation of two nonempty sets $A \subset S$ and $B = S \setminus A$ by an affine hyperplane H . Throughout this paper we assume that A, B are nonempty disjoint sets and we denote $S = A \cup B$. H may be defined by a linear form with irrational coefficients.

We allow H to contain points of both A and B . However on H we are again concerned on separation of A and B by an affine space of codimension 1 within H . Therefore we make the following definition.

Definition 2.1. Let L be an affine subspace in $\mathbb{R}^d$ of dimension l . We call an affine subspace $P \subset L$ of dimension $l - 1$ a *hyperplane* in L . We say that

P separates A and B in L if there exist two disjoint connected components R^+ and R^- of $L \setminus P$ such that

$$R^+ \cup R^- = L \setminus P, \quad R^+ \cap S \subset A, \quad R^- \cap S \subset B.$$

Example 2.2. We consider the case when $A = \{0, 1, 2\}$, $B = \{-1, -2\}$. Let us take the rays $\mathbb{R}_{>0}$ and $\mathbb{R}_{<0}$ as R^+ and R^- , respectively. Then $P = \mathbb{R} \setminus (R^+ \cup R^-)$ has only the point $0 \in A$. In this case, $S \cap R^+ = \{1, 2\} \subset A$ and $S \cap R^- = \{-1, -2\} \subset B$. Hence $P = \{0\}$ separates A and B in $\mathbb{R}$.

Definition 2.3 (Separation by a sequence of affine hyperplanes, Condition H). Let H_i ($i \geq 0$) be an i -dimensional affine subspaces of $\mathbb{R}^d$ and $H = \{H_k, \dots, H_d\}$ ($k \geq 0$) be a sequence of affine hyperplanes such that $H_{i-1} \subset H_i$ ($i = k+1, \dots, d$) and $H_d = \mathbb{R}^d$. A and B are separated by H , if H satisfies

- (i) $H_k \cap S \subset A$ or $H_k \cap S \subset B$, and
- (ii) for $k+1 \leq i \leq d$, H_{i-1} separates A and B in H_i .

We say that A and B satisfy *Condition H* if A and B are separated by some $H = \{H_k, \dots, H_d\}$.

We consider two examples of separation of $A \subset \mathbb{Z}^2$ and $B = \mathbb{Z}^2 \setminus A$.

Example 2.4. Let H_1 be the line defined by $x_2 = \varpi x_1$ for some irrational number ϖ , $A = \{(x_1, x_2) \in \mathbb{Z}^2 \mid x_2 \geq \varpi x_1\}$, and $H_2 = \mathbb{R}^2$. Then $H = \{H_1, H_2\}$ separates A and $B = \mathbb{Z}^2 \setminus A$. (Moreover, for any t , $H' = \{(t, \varpi t)\}, H_1, H_2\}$ separates A and B .)

Remark 2.5. For $S = \mathbb{Z}^d$ there may be no rational vector $p \in \mathbb{Z}^d$ such that a hyperplane $H = \{x \in \mathbb{R}^d \mid p_1 x_1 + \dots + p_d x_d = b\}$ separates A and B , because $\mathbb{Z}^d$ is unbounded.

Example 2.6. Let $A = \{(x_1, x_2) \in \mathbb{Z}^2 \mid x_1 > 0\} \cup \{(0, x_2) \in \mathbb{Z}^2 \mid x_2 \geq 0\}$ as in Figure 1. If we set

$$H_0 = \{(0, 0)\}, \quad H_1 = \{(0, x_2) \mid x_2 \in \mathbb{R}\}, \quad H_2 = \mathbb{R}^2,$$

then $H = \{H_0, H_1, H_2\}$ separates A and $B = \mathbb{Z}^2 \setminus A$. Moreover, for each $t \in [-1, 0]$, $\{(0, t)\}, H_1, H_2\}$ separates A and B .

Next we consider the following condition.




Figure 1: Example 2.6

Definition 2.7 (Ray condition, Condition R). We say that A and B satisfy *Condition R* if for each line L such that $A \cap L \neq \emptyset$ and $B \cap L \neq \emptyset$, there exists a ray $L' \subset L$ such that $A \cap L = L' \cap S$.

Remark 2.8. Condition R is equivalent to the following: for each line L such that $A \cap L \neq \emptyset$ and $B \cap L \neq \emptyset$, there exists $p \in L$ such that $\{p\}$ separates A and B in L .

Furthermore we consider the following condition.

Definition 2.9 (Parallelogram condition, Condition P). We say that A, B satisfy *k-parallelogram condition* if

$$a_1, a_2, \dots, a_{k'} \in A, b_1, b_2, \dots, b_{k'} \in B \Rightarrow \sum_{i=1}^{k'} a_i \neq \sum_{i=1}^{k'} b_i$$

for all $k' \leq k$. We call the 2-parallelogram condition *Condition P*.

Remark 2.10. By definition, the k -parallelogram condition implies the $(k-1)$ -parallelogram condition.

When we consider Condition P, we may have $a_1 = a_2$. Then the condition says that no point of A is the mid-point of two points in B . Condition P was considered in Hara et al. [2] in a statistical problem. They showed that when $S = A \cup B$ is a 2-dimensional rectangle, and if A and B are “monotone” and satisfy Condition P, then there exists a line separating A and B (see Appendix E in Hara et al. [2]).

We state the following basic fact on implications of the above conditions.

Lemma 2.11. *Condition H implies Condition P.*

Proof. Consider the case that at least one point, say a_1 , among $a_1, a_2 \in A, b_1, b_2 \in B$ belongs to an open half-space on one side of H_{d-1} . Then $(a_1 + a_2)/2$ also belongs to the open half-space and can not be equal to $(b_1 + b_2)/2$. If all of a_1, a_2, b_1, b_2 belong to H_{d-1} , then we can use the induction on dimension. $\square$

By the same proof as in Lemma 2.11, we have the following.

Lemma 2.12. *Condition H implies the k -parallelogram condition for every $k \geq 2$.*

So far we have presented conditions concerning two sets A and B . Now we give a condition of discrete convexity of a single S , which is also needed for relating Condition R to Condition P.

Definition 2.13 (k -convexity). $S \subset \mathbb{Z}^d$ is k -convex if S satisfies the following:

$$a_1, \dots, a_{k+1} \in S \Rightarrow \text{conv} \{ a_1, \dots, a_{k+1} \} \cap \mathbb{Z}^d \subset S,$$

where $\text{conv} \{ a_1, \dots, a_{k+1} \}$ denotes the convex hull of $a_1, \dots, a_{k+1}$ in $\mathbb{R}^d$.

Remark 2.14. If $S \subset \mathbb{Z}^d$ is k -convex, then S is $(k-1)$ -convex.

Lemma 2.15. *If $S = A \cup B$ is 1-convex, then Condition P implies Condition R.*

Proof. If $A \cap L \neq \emptyset, B \cap L \neq \emptyset$, then we can find $a \in A$ and $b \in B$ lying next to each other on the line L by 1-convexity of S . Assume that a is on the right side of b , and let $L' \subset L$ denote the ray extending from a not containing b . If there exists a member of B in $L' \cap \mathbb{Z}^d$, then we find the nearest point to b in B , denoted by b' . Let a' denote the next left point of b' in $L' \cap \mathbb{Z}^d$. Then we have $a' \in A$ and $a + a' = b + b'$, a contradiction to Condition P. Thus $L' \cap \mathbb{Z}^d \subset A$. Also we have $(L \setminus L') \cap \mathbb{Z}^d \subset B$. Then it follows that $L' \cap \mathbb{Z}^d = A \cap L$. $\square$

Lemma 2.16. *If S is 1-convex, then*

$$\text{Condition H} \Rightarrow \text{Condition P} \Rightarrow \text{Condition R}.$$

Proof. The lemma follows from Lemmas 2.11 and 2.15. $\square$

In Section 4 we consider the case when S is a hole free set or an integrally convex set. We recall its definition and basic properties ([7, Section 3.4]). See also [1].

Definition 2.17. $S \subset \mathbb{Z}^d$ is *integrally convex* if

$$x \in \text{conv}(S) \Rightarrow x \in \text{conv}(S \cap N(x)),$$

where $N(x) = \{ y \in \mathbb{Z}^d \mid |x_i - y_i| < 1 \text{ for } i = 1, \dots, d \}$ is the integral neighborhood of $x \in \mathbb{R}^d$.

This definition is also written as follows.

Proposition 2.18. $S \subset \mathbb{Z}^d$ is *integrally convex* if and only if

$$\text{conv}(S \cap N(x)) = \text{conv}(S) \cap \text{conv}(N(x)) \quad (\forall x \in \mathbb{R}^d).$$

Definition 2.19. $S \subset \mathbb{Z}^d$ is *hole free* if $S = \text{conv}(S) \cap \mathbb{Z}^d$.

Proposition 2.20. An *integrally convex* set is *hole free*.

Remark 2.21. For $S \subset \mathbb{Z}^d$, S is d -convex if and only if S is hole free.

3 Separation of the whole integer lattice

In this section, we consider the case $S = \mathbb{Z}^d$. We prove the following theorem.

Theorem 3.1. Let $S = \mathbb{Z}^d$. Then for $A \subset S$ and $B = S \setminus A$, Conditions H , R , P are all equivalent.

By Lemma 2.16 it suffices to prove Condition $R \Rightarrow$ Condition H . We give the proof in a series of lemmas and the following key proposition.

Proposition 3.2. Let $l < n \leq d$ and $S = \mathbb{Z}^d \cap F$ for some affine subspace $F \subset \mathbb{R}^d$ of dimension n . Assume A and B satisfy Condition R . If there exist affine subspaces $P \subset L$ of dimension $(l - 1)$ and l such that P separates A and B in L , then there exist affine subspaces $P' \subset L'$ of dimension l and $(l + 1)$ such that P' separates A and B in L' .

For readability we shall prove this proposition later. It implies the following two lemmas, which are sufficient to prove Theorem 3.1.

Lemma 3.3. *Let $S = \mathbb{Z}^d \cap F$ for some affine subspace $F \subset \mathbb{R}^d$. If A and B satisfy Condition R, then there exists a hyperplane $H \subset F$ such that H separates A and B in F .*

Proof. Let $a \in A$ and $b \in B$. For the line L through a and b , some point $p \in L$ separates A and B in L since A and B satisfy Condition R. Applying Proposition 3.2 inductively, we have the lemma. $\square$

Lemma 3.4. *Let $S = \mathbb{Z}^d \cap F$ for some affine subspace $F \subset \mathbb{R}^d$ of dimension n . If A and B satisfy Condition R, then Condition H holds.*

Proof. By Lemma 3.3, there exists a hyperplane H_{n-1} in $H_n = F$ such that H_{n-1} separates A and B in H_n . Applying Lemma 3.3 recursively until $H_i \subset A$ or $H_i \subset B$, we have a sequence of hyperplanes which separate A and B . $\square$

Proof of Theorem 3.1. Take $F = \mathbb{R}^d$ in Lemma 3.4. $\square$

3.1 Proof of Proposition 3.2

It remains to prove Proposition 3.2. The rest of this section is devoted to its proof in a series of lemmas. For $1 \leq i \leq d$, let

$$R_i(\theta) = \{ (x_1, \dots, x_{i-1}, t \cos \theta, t \sin \theta, 0, \dots, 0) \in \mathbb{R}^d \mid x_1, \dots, x_{i-1} \in \mathbb{R}, t > 0 \},$$

$$H_i = \{ (x_1, \dots, x_i, 0, \dots, 0) \in \mathbb{R}^d \mid x_1, \dots, x_i \in \mathbb{R} \},$$

$R_i^+ = R_i(0)$ and $R_i^- = R_i(\pi)$. Assume $\varphi \in \text{GL}(\mathbb{R}^d)$ and $\alpha \in \mathbb{R}^d$ satisfy the following: $P = \varphi(H_{l-1}) + \alpha$, $L = \varphi(H_l) + \alpha$, $F = \varphi(H_n) + \alpha$, $R^+ = \varphi(R_l^+) + \alpha$, $R^- = \varphi(R_l^-) + \alpha$, $R^+ \cap S \subset A$ and $R^- \cap S \subset B$.

Lemma 3.5. *Let $a \in R^+ \cap S = (\varphi(R_l^+) + \alpha) \cap S$. There exist θ'_a, θ''_a , and C_a such that*

$$\theta'_a - \theta''_a \geq \pi, \quad C_a = \bigcup_{\theta''_a \leq \theta \leq \theta'_a} (\varphi(R_l(\theta)) + a), \quad \text{Int}(C_a) \cap S \subset A,$$

where $\text{Int}(X)$ denotes the relative interior of $X \subset \mathbb{R}^d$.

Proof. Define θ'_a, θ''_a as

$$\begin{aligned}\theta'_a &= \sup \{ \theta' \mid \forall \theta \in [0, \theta'], (\varphi(R_l(\theta)) + a) \cap S \subset A \}, \\ \theta''_a &= \inf \{ \theta'' \mid \forall \theta \in [\theta'', 0], (\varphi(R_l(\theta)) + a) \cap S \subset A \}.\end{aligned}$$

Since $(\varphi(R_l^+) + a) \subset (\varphi(R_l^+) + \alpha) \subset A$ and $\emptyset \neq (\varphi(R_l^-) + \alpha) \subset B$, it follows that $-\pi \leq \theta''_a \leq 0 \leq \theta'_a \leq \pi$. Let $C_a = \bigcup_{\theta''_a \leq \theta \leq \theta'_a} (\varphi(R_l(\theta)) + a)$. Then we have $\text{Int}(C_a) \cap S \subset A$.

The proof is completed by showing that $\theta'_a - \theta''_a \geq \pi$. For convenience, let $a = 0$. If $\theta'_a - \theta''_a < \pi$, then there exist θ', θ'' such that

$$\begin{aligned}\theta'' &\leq \theta''_a \leq \theta'_a \leq \theta' < \theta'' + \pi, \\ \varphi(R_l(\theta'')) \cap B &\neq \emptyset, \\ \varphi(R_l(\theta')) \cap B &\neq \emptyset.\end{aligned}$$

Let $b' \in \varphi(R_l(\theta')) \cap B, b'' \in \varphi(R_l(\theta'')) \cap B$. Since $b', b'' \in \mathbb{Z}^d$, there exist $t', t'' \in \mathbb{Z}$ such that

$$\frac{t'b' + t''b''}{t' + t''} \in \varphi(R_l^+).$$

Let $q' = (t' + t'')b', q'' = (t' + t'')b''$, and $p = (t'q' + t''q'')/(t' + t'') = t'b' + t''b''$. Then $q', q'' \in B$, and $p \in (\varphi(R_l^+) + a) \cap \mathbb{Z}^d \subset A$. However the three points p, q', q'' contradict Condition R. $\square$

Lemma 3.6. *Let $b \in R^- \cap S = (\varphi(R_l^-) + \alpha) \cap S$. There exist θ'_b, θ''_b , and C_b such that*

$$\theta'_b - \theta''_b \geq \pi, \quad C_b = \bigcup_{\theta''_b \leq \theta \leq \theta'_b} (\varphi(R_l(\theta)) + b), \quad \text{Int}(C_b) \cap \mathbb{Z}^d \subset B.$$

Proof. The proof is the same as that of Lemma 3.5. $\square$

Since $\text{Int}(C_a) \cap S \subset A, \text{Int}(C_b) \cap S \subset B$ and $A \cap B = \emptyset$, it immediately follows that $\theta'_a - \theta''_a = \pi$. Since

$$C_a = \bigcup_{\theta''_a \leq \theta \leq \theta'_a} (\varphi(R_l(\theta)) + a) = \varphi \left(\bigcup_{\theta''_a \leq \theta \leq \theta'_a} R_l(\theta) \right) + a,$$

there exists $\psi_a \in \text{GL}(\mathbb{R}^d)$ such that $\psi_a(R_{l+1}^+) + a = \text{Int}(C_a)$.

Let us fix $a \in R^+ \cap S$ and $\psi = \psi_a$. Let $L' = \psi(H_{l+1}) + a$ and $C'_{a'} = C_a - a + a' \subset L'$ for $a' \in L' \cap A$.

Lemma 3.7. *For $a' \in L' \cap A$, $\text{Int}(C'_{a'}) \cap S \subset A$.*

Proof. By Lemma 3.5, $\text{Int}(C_a) \subset A$. If $a' \in C_a$, then $C'_{a'} \subset C_a$, which implies $\text{Int}(C'_{a'}) \subset \text{Int}(C_a) \subset A$. Let us consider the case when $a' \notin C_a$. For each $x \in \text{Int}(C'_{a'}) \cap S$, there exists some point $q \in C_a \cap S \subset A$ on the ray from a to x . Since we assume Condition R, x is in A . $\square$

Let

$$\begin{aligned} R'_+ &= \bigcup_{a' \in L' \cap A} \text{Int}(C'_{a'}) = \bigcup_{a' \in L' \cap A} \text{Int}(C_a) - a + a' \\ &= \bigcup_{a' \in L' \cap A} (\psi(R_{l+1}^+) + a'). \end{aligned}$$

Fix $\beta \in \partial R'_+ = \overline{R'_+} \setminus \text{Int}(R'_+)$. Then we obtain

$$R'_+ = \psi(R_{l+1}^+) + \beta, \quad L' \setminus \overline{R'_+} = \psi(H_{l+1}) \setminus \overline{R'_+} = \psi_a(R_{l+1}^-) + \beta.$$

Let $R'_- = \psi_a(R_{l+1}^-) + \beta$ and $P' = L' \setminus (R'_+ \cup R'_-)$. Then we turn to show that P' separates A and B in L' .

Lemma 3.8. $\emptyset \neq R'_- \cap S \subset B$.

Proof. Since $A \cap L' \subset \overline{R'_+}$, it follows that $S \cap (L' \setminus \overline{R'_+}) \subset B$, which implies $S \cap R'_- \subset B$. $\square$

Lemma 3.9. $\emptyset \neq R'_+ \cap S \subset A$.

Proof. For $x \in R'_+$, there exists $a' \in L'$ such that $x \in \text{Int}(C'_{a'})$, which implies $x \in A$. $\square$

Now we have Proposition 3.2. Note that in the above proofs we described how to construct a sequence of separating affine subspaces.

4 Separation of a finite set in dimension two

In this section, we consider the case when $S \subset \mathbb{Z}^d$ is finite. In the case $d = 2$ simple separation results hold as shown in the following theorems.

Theorem 4.1. *Let $S \subset \mathbb{Z}^2$ be a finite integrally convex set. Then for $A \subset S$ and $B = S \setminus A$, Conditions H and P are equivalent.*

Theorem 4.2. *Let $S \subset \mathbb{Z}^2$ be a finite hole free set. Then for $A \subset S$ and $B = S \setminus A$, Condition H and the 3-parallelogram condition are equivalent.*

We will prove these theorems in later subsections by using the following lemma.

Lemma 4.3. *Let $S \subset \mathbb{Z}^2$ be a finite hole free set. If A and B satisfy Condition P, $A \setminus \text{conv}(B) \neq \emptyset$ and $B \setminus \text{conv}(A) \neq \emptyset$, then Condition H holds.*

We show the following lemma before the proof of Lemma 4.3.

Lemma 4.4. *Let $S \subset \mathbb{Z}^2$ be a finite hole free set. If A and B satisfy Condition P, then $\text{conv}\{a_1, a_2\} \cap \text{conv}\{b_1, b_2\} = \emptyset$ for all $a_1, a_2 \in A$ and $b_1, b_2 \in B$.*

Proof. Let us assume that for some $a_1, a_2 \in A$ and $b_1, b_2 \in B$, $\text{conv}\{a_1, a_2\} \cap \text{conv}\{b_1, b_2\} \neq \emptyset$. Consider a_1, a_2, b_1, b_2 such that $\text{conv}\{a_1, a_2, b_1, b_2\} \cap \mathbb{Z}^2$ is minimal with respect to inclusion. Thanks to the hole freeness of S , we can suppose that $\text{conv}\{a_1, a_2, b_1, b_2\} \cap \mathbb{Z}^2 = \{a_1, a_2, b_1, b_2\}$ without loss of generality.

Let $\delta = a_1 - a_2$ and $c = a_1 + a_2 - b_1$. Also let $c_+ = c + \delta, c_- = c - \delta$. Since the area of the triangle $\text{conv}\{a_1, a_2, b_1\}$ is equal to that of the triangle $\text{conv}\{a_1, a_2, b_2\}$, b_2 has to be on the line parallel to $\text{conv}\{a_1, a_2\}$ through c . Also $\text{conv}\{a_1, a_2, b_1, b_2\} \cap \mathbb{Z}^2 = \{a_1, a_2, b_1, b_2\}$ implies $b_2 \in \text{conv}\{c_+, c_-\}$. Then since $\text{conv}\{a_1, a_2\} \cap \mathbb{Z}^2 = \{a_1, a_2\}$, we have

$$\text{conv}\{c_+, c_-\} \cap \mathbb{Z}^2 = \{c, c_+, c_-\}.$$

Thus $b' \in \{c, c_+, c_-\}$. However in any case we have a contradiction to Condition P. $\square$

Proof of Lemma 4.3. If $\text{conv}(A) \cap \text{conv}(B) = \emptyset$, then Condition H holds by the usual hyperplane separation theorem. Therefore it suffices to show that $\text{conv}(A) \cap \text{conv}(B) \neq \emptyset$ leads to a contradiction.

Firstly, recall that the convex hull of a finite set of points is equal to the union of simplices spanned by a finite set of the points [11, Thm.2.15]. Therefore, if $\text{conv}(A) \cap \text{conv}(B) \neq \emptyset$, there exist $a_1, a_2, a_3 \in A$ and $b_1, b_2, b_3 \in B$ such that $\text{conv}\{a_1, a_2, a_3\} \cap \text{conv}\{b_1, b_2, b_3\} \neq \emptyset$. By Lemma 4.4, it suffices to show that for some $a, a' \in A$ and $b, b' \in B$, $\text{conv}\{a, a'\}$ and $\text{conv}\{b, b'\}$ intersect.

Let $A' = \{a_1, a_2, a_3\}$ and $B' = \{b_1, b_2, b_3\}$. As the first case suppose that some $b \in B'$ is in $\text{conv}(A')$. Then since $B \setminus \text{conv}(A) \neq \emptyset$, there exists $b' \in B$ such that $b' \notin \text{conv}(A')$. In this case, clearly the line segment $\text{conv}\{b, b'\}$ crosses an edge of the triangle $\text{conv}(A')$.

As the second case suppose that $B' \cap \text{conv}(A') = \emptyset$ and $A' \cap \text{conv}(B') = \emptyset$. The intersection of two triangles $\text{conv}(A') \cap \text{conv}(B')$ is a polygon. Consider a vertex v of the polygon. Since $v \in \text{conv}(A') \cap \text{conv}(B')$, v is not a member of $A' \cup B'$. Now each triangle is defined by three inequalities and the intersection of the two triangles is defined by effective inequalities of those six. Hence v is the intersection of two edges among these six edges. If v is the intersection of two edges of one triangle, then it reduces to the first case. Hence v is the intersection of one edge of $\text{conv}(A')$ and another edge of $\text{conv}(B')$. $\square$

In the finite case, note that Condition R is not equivalent to Conditions P and H.

Example 4.5 (A counterexample to $R \Rightarrow H$). When $A = \{(0, 0), (1, 1)\}$ and $B = \{(1, 0), (0, 1)\}$, $S = A \cup B$ is integrally convex. In this case, both A and B satisfy Condition R. However we easily find that Conditions H and P do not hold.

In the case $d = 2$, we have Theorems 4.1 and 4.2. We want to consider more general cases or higher dimensions. However, there are some counterexamples.

Firstly we consider Theorem 4.1. For general sets $S \subset \mathbb{Z}^2$, the equivalence of Theorem 4.1 is violated as shown in the following examples.

Example 4.6 (Counterexamples to $P \Rightarrow H$). We show some counterexamples in Figure 2. The points of A are denoted by black circles, and those of B by white circles. In these cases, Condition P holds (moreover Condition R also holds from Lemma 2.16). However, A and B are not separated.

We next consider the case $S \subset \mathbb{Z}^d$ for $d \geq 3$. In this case, by examples we show that Conditions H, P and R are not equivalent.

Example 4.7. Consider

$$S = \{ (x_1, x_2, x_3) \in \mathbb{Z}^3 \mid 0 \leq x_1 \leq 5, 0 \leq x_2 \leq 4, 0 \leq x_3 \leq 3 \}.$$

Let v_1, v_2, v_3 denote $(5, 0, 0), (0, 4, 0), (0, 0, 3)$ and $S' = \text{conv}\{0, v_1, v_2, v_3\} \cap \mathbb{Z}^3$ as in Figure 3, $A = S' \setminus \{(2, 1, 1)\}$ and $B = S \setminus A$. Then $\text{conv}(A) \cap \mathbb{Z}^3 = S'$ and it is clear that there is no hyperplane separating A and B . However we can easily check that Condition P holds.




Figure 2: Some counterexamples to $P \Rightarrow H$ in $\mathbb{Z}^2$




Figure 3: A counterexample in $\mathbb{Z}^3$

In this example, we see that the set A is the smallest 1-convex set containing $\{0, v_1, v_2, v_3\}$, which we call the 1-convex hull of $\{0, v_1, v_2, v_3\}$. Similarly define the k -convex hull of S as the smallest k -convex set containing S . In this case, since we have

$$\frac{1}{3}(5, 0, 0) + \frac{1}{3}(0, 0, 3) + \frac{1}{3}(1, 3, 0) = (2, 1, 1),$$

S is 1-convex but it is not 2-convex (hence not 3-convex). Therefore the 2-convex hull of $\{0, v_1, v_2, v_3\}$ is the same as S' .

Then we are interested in the following example which shows that the $(d - 1)$ -convex hull is not necessarily d -convex.

Example 4.8. Let v_1, v_2, v_3 denote the points $(13, 0, 0), (0, 7, 0), (0, 0, 4)$ and $S' = \text{conv}\{0, v_1, v_2, v_3\} \cap \mathbb{Z}^3$. Then by some detailed calculation involving inner products, it can be shown that there exists no $a_1, a_2, a_3 \in S'$ such that

$$(6, 2, 1) \in \text{conv}\{a_1, a_2\} \quad \text{or} \quad (6, 2, 1) \in \text{conv}\{a_1, a_2, a_3\}.$$

Hence, the 2-convex hull of $\{0, v_1, v_2, v_3\}$ is not 3-convex.

This example shows that the equivalence of Theorem 4.2 is also violated for $d \geq 3$.

Thus we are interested in generalizations of Theorems 4.1 and 4.2 using the k -parallelogram condition where $k \geq d$. However the following counterexamples suggest that generalizations of Theorem 4.2 and Lemma 4.3 are difficult for $d \geq 3$.

Example 4.9. (An example based on a terminal Fano polytope in $\mathbb{R}^3$)

Let v_1, v_2, v_3, v_4 denote the points $(1, 0, 0), (0, 1, 0), (1, 1, 2), (-1, -1, -1)$ and $S' = \text{conv}\{v_1, v_2, v_3, v_4\}$. Then since

$$v_1 + v_2 + v_3 + 2v_4 = 0,$$

we have $0 \in S'$. Let $A = S' \setminus \{0\}$ and $B = \{0\}$. Since S' has no lattice points other than $v_1, \dots, v_4, 0$, we have the 4-parallelogram condition. However we can not separate A and B .

Example 4.10. Let a_1, a_2, a_3 denote the points $(1, 0, 0), (0, 1, 0), (0, 0, 1)$ and b_1, b_2 denote $(0, 0, 0), (1, 1, 2)$. Let $A = \{a_1, a_2, a_3\}$, $B = \{b_1, b_2\}$. Then $S = A \cup B$ is hole free and $A \cap \text{conv}(B) = B \cap \text{conv}(A) = \emptyset$. We clearly have the 3-parallelogram condition, but not Condition H.

Despite the above counterexamples, we conjecture that a generalization of Theorem 4.1 to $d \geq 3$ holds (see Section 5).

4.1 Proof of Theorem 4.1

For proving Theorem 4.1 we show the following basic lemmas on an integrally convex set.

Lemma 4.11. *For a line L in $\mathbb{R}^d$, if $L \cap \mathbb{Z}^d$ is integrally convex, then L is of the form*

$$L = \{x_0 + tp \mid t \in \mathbb{R}\} \tag{1}$$

for some $x_0 \in \mathbb{Z}^d$ and $p \in \{-1, 0, 1\}^d \setminus \{0\}$.

Proof. Let $I = L \cap \mathbb{Z}^d$. For an integer point $x_0 \in L \cap \mathbb{Z}^d$, choose $\alpha \in (-1, 1)^d \setminus \{0\}$ such that $x_0 + \alpha \in L \setminus I$. We denote $x = x_0 + \alpha$. Consider the intersection of I and the integral neighborhood $N(x)$. Since $L = \text{conv}(I)$ is a line, $I \cap N(x)$ contains two points, and one of them is x_0 . Thus another one has to be $x_0 + p$ where $p \in \{-1, 0, 1\}^d \setminus \{0\}$. Then we have the lemma. $\square$

Lemma 4.12. *The faces of a finite integrally convex set in $\mathbb{R}^d$ are integrally convex.*

Proof. Let $S \subset \mathbb{Z}^d$ be a finite integrally convex set. From Proposition 2.18, for $x \in \mathbb{R}^d$, we can consider $\text{conv}(S \cap N(x))$ as $\text{conv}(S)$ restricted to $\text{conv}(N(x))$. For a face F of $\text{conv}(S)$ and $x \in F$, let $F_x = F \cap \text{conv}(N(x))$. Since F_x is a face of $\text{conv}(S) \cap \text{conv}(N(x))$, it suffices to show that $F_x \cap \mathbb{Z}^d$ is integrally convex.

Let $P \subset \mathbb{R}^d$ be a polytope, $\text{vert}(P)$ be the set of all vertices of P , and E be a face of P . Then

$$P = \text{conv}(\text{vert}(P)), \quad (2)$$

$$\text{vert}(E) = E \cap \text{vert}(P). \quad (3)$$

Since $\text{conv}(S \cap N(x))$ is a 0/1-polytope, we have $\text{vert}(\text{conv}(S \cap N(x))) = S \cap N(x)$. Thus by (3) we have

$$\begin{aligned} \text{vert}(F_x) &= F_x \cap \text{vert}(\text{conv}(S \cap N(x))) \\ &= F_x \cap S \cap N(x) = F_x \cap N(x). \end{aligned}$$

Therefore, for any $y \in F_x$,

$$\begin{aligned} \text{conv}(\text{vert}(F_x) \cap N(y)) &= \text{conv}(F_x \cap N(x) \cap N(y)) \\ &= \text{conv}(F_x \cap N(y)). \end{aligned} \quad (4)$$

Since $F_x \cap \text{conv}(N(y))$ is a face of F_x , we have

$$\begin{aligned} \text{vert}(F_x \cap \text{conv}(N(y))) &= F_x \cap \text{conv}(N(y)) \cap \text{vert}(F_x) \\ &= \text{vert}(F_x) \cap N(y). \end{aligned} \quad (5)$$

Thus combining the equations (2), (4) and (5), we see that $F_x \cap \mathbb{Z}^d$ is integrally convex. $\square$

From these lemmas, as an edge of an integrally convex set in $\mathbb{Z}^2$ we only need to consider lines of the form (1). We now finish our proof of Theorem 4.1. In view of Lemma 4.3, suppose that $B \subset \text{conv}(A)$ or $A \subset \text{conv}(B)$. Without loss of generality let $B \subset \text{conv}(A)$. Then

$$S = A \cup B \subset \text{conv}(A) \cup B = \text{conv}(A).$$

Since $S \supset A$, we have $\text{conv}(S) = \text{conv}(A)$. Thus all vertices of $\text{conv}(S)$ belong to A . This implies that the integer points of the boundary of $\text{conv}(S)$ also belong to A by Condition P. Let $b \in B$. Above lemmas show that as we move from b in a direction parallel to either axis, we reach an integer point of an edge of S , which belongs to A . If we move to the opposite direction of b we again reach an integer point of A . This contradicts Condition P.

4.2 Proof of Theorem 4.2

Finally we give a proof of Theorem 4.2. When we look at counterexamples in Figure 2 again, we notice that the centroid of black circles is also a centroid of white circles. Therefore these counterexamples are impossible if we additionally impose the 3-parallelogram condition.

Let $S \subset \mathbb{Z}^2$ be hole free. Since Lemma 4.3 holds, we only need to prove that the 3-parallelogram condition implies $A \cap \text{conv}(B) = \emptyset$ and $\text{conv}(A) \cap B = \emptyset$. We prove it by contradiction. Note that we only need to consider minimal triangles in $\text{conv}(A)$, i.e., triangles with no points from A except the vertices a_1, a_2, a_3 . Moreover, each face of this triangle has no integer points except them since we have the 2-parallelogram condition.

Theorem 4.2 is a consequence of the following lemma.

Lemma 4.13. *Let $A = \{a_1, a_2, a_3\}$. Assume that $\text{conv}\{a_i, a_j\} \cap \mathbb{Z}^2 = \{a_i, a_j\}$ for any $i, j \in \{1, 2, 3\}$ and $B = (\text{conv}(A) \cap \mathbb{Z}^2) \setminus A$ is not empty. Then there exist $b_1, b_2, b_3 \in B$ such that $a_1 + a_2 + a_3 = b_1 + b_2 + b_3$.*

Proof. We can assume that $a_3 = 0$ without loss of generality. Let us fix $a_1^\perp$, $a_2^\perp$ and $a_{12}^\perp$ such that

$$\begin{aligned} \langle a_1, a_1^\perp \rangle &= 0, & \langle a_2, a_2^\perp \rangle &= 0, & \langle a_1 - a_2, a_{12}^\perp \rangle &= 0, \\ \langle a_2, a_1^\perp \rangle &> 0, & \langle a_1, a_2^\perp \rangle &> 0, & \langle -a_1, a_{12}^\perp \rangle &= \langle -a_2, a_{12}^\perp \rangle > 0. \end{aligned}$$

Then, for $x \in \mathbb{R}^2$,

$$x \in \text{conv}(A) \iff \langle x, a_1^\perp \rangle \geq 0, \langle x, a_2^\perp \rangle \geq 0, \langle x - a_1, a_{12}^\perp \rangle \geq 0.$$

By definition it also follows that $\langle b - a_2, a_1^\perp \rangle < 0$, $\langle b - a_1, a_2^\perp \rangle < 0$ and




Figure 4: A minimal triangle

$\langle b, a_{12}^\perp \rangle < 0$ for all $b \in B$. Let

$$\begin{aligned} b_1 &= \operatorname{ArgMax}_{b \in B} \langle b - a_1, a_2^\perp \rangle, \\ b_2 &= \operatorname{ArgMax}_{b \in B} \langle b - a_2, a_1^\perp \rangle, \\ b_3 &= a_1 + a_2 - b_1 - b_2. \end{aligned}$$

We show that $b_3 \in B$, which concludes the lemma. Let $c = b_1 + b_2 - a_1$. Since $\langle c - a_2, a_1^\perp \rangle = \langle b_1 - a_2, a_1^\perp \rangle + \langle b_2, a_1^\perp \rangle > \langle b_1 - a_2, a_1^\perp \rangle$, by the definition of b_2 , we get $c \notin \operatorname{conv}(A)$. By direct calculation we obtain

$$\begin{aligned} \langle c, a_1^\perp \rangle &= \langle b_1, a_1^\perp \rangle + \langle b_2, a_1^\perp \rangle > 0, \\ \langle c - a_1, a_{12}^\perp \rangle &= \langle b_1 - a_1, a_{12}^\perp \rangle + \langle b_2 - a_1, a_{12}^\perp \rangle > 0, \end{aligned}$$

which implies $\langle c, a_2^\perp \rangle < 0$. Since $b_3 = a_2 - c$, $\langle b_3, a_2^\perp \rangle = \langle a_2 - c, a_2^\perp \rangle = -\langle c, a_2^\perp \rangle > 0$. Similarly we also obtain $c' = b_1 + b_2 - a_2 \notin \operatorname{conv}(A)$ and $\langle b_3, a_1^\perp \rangle > 0$. Since $\langle b_3 - a_1, a_{12}^\perp \rangle = \langle b_3, a_{12}^\perp \rangle + \langle -a_1, a_{12}^\perp \rangle > 0$, we have $b_3 \in \operatorname{Int}(\operatorname{conv}(A))$, which implies $b_3 \in B$. $\square$

5 Conclusions and open problems

In this paper, we have proved the equivalence of Conditions H and P when $S = A \cup B$ is the whole $\mathbb{Z}^d$ or a 2-dimensional finite integrally convex set.

However we have shown that it does not hold in d -dimensional finite sets for $d \geq 3$. For $S = \mathbb{Z}^d$ Condition R is also equivalent to Conditions H and P, but they are not equivalent in the finite case. It is not clear what causes this difference. We have also shown the equivalence of Condition H and the 3-parallelogram condition for finite hole free $S \subset \mathbb{Z}^2$.

For $d \geq 3$ and finite S , we would like to obtain a useful condition which is equivalent to Condition H. Such a condition is relevant for extending the result of [2] to higher dimensions. From various counterexamples in Section 4, extension of our results for $d = 2$ to higher dimensions may be difficult to prove. In fact, in Examples 4.9 and 4.10 we see that a natural generalization of Theorem 4.2 does not hold. Nevertheless, we conjecture that a generalization of Theorem 4.1 holds in higher dimensions.

Conjecture 5.1 (Analogue of Theorem 4.1). *If S is a finite integrally convex set in $\mathbb{R}^d$ and A, B satisfy the d -parallelogram condition, then Condition H holds.*

Another open problems concern the k -convexity of a set $A \in \mathbb{Z}^d$. The k -convex hull of A is a subset of the $(k + 1)$ -convex hull of A . A hole in $\text{conv}(A) \cap \mathbb{Z}^d$ can be classified by the first k such that it belongs to the k -convex hull of A . For example, in Example 4.8 with $A = \{(0, 0, 0), (13, 0, 0), (0, 7, 0), (0, 0, 4)\}$ by computer search we found that the difference between the 2-convex hull and the 1-convex hull consists of a single point $(4, 3, 1)$. $(6, 2, 1)$ discussed in Example 4.8 the difference between the 3-convex hull and the 2-convex hull. It is of interest to classify holes by the k -convexity and give conditions on existence or non-existence of holes according to this classification.

Acknowledgment. We are grateful to Kazuo Murota for very helpful comments.

References

- [1] P. Favati and F. Tardella, Convexity in nonlinear integer programming, *Ricerca Operativa* **53** (1990), 3–44.
- [2] H. Hara, A. Takemura and R. Yoshida, On connectivity of fibers with positive marginals in multiple logistic regression, *Journal of Multivariate Analysis* **101** (2010), 909–925.

- [3] R. Hemmecke, A. Takemura and R. Yoshida, Computing holes in semi-groups and its applications to transportation problems, *Contributions to Discrete Mathematics* **4** (2009), 81–91.
- [4] A. M. Kasprzyk, Toric Fano threefolds with terminal singularities, *Tohoku Mathematical Journal (2)* **58** no. 1 (2006), 101–121.
- [5] A. M. Kasprzyk, Canonical toric Fano threefolds, preprint (2008). [arXiv:0806.2604v2](#). To appear in *Canadian Journal of Mathematics*.
- [6] K. Murota, Discrete convex analysis, *Mathematical Programming* **83** (1998), 313–371.
- [7] K. Murota, *Discrete Convex Analysis*, Monographs on Discrete Mathematics and Applications, SIAM, Philadelphia, 2003.
- [8] K. Murota, Recent developments in discrete convex analysis, in: W. Cook, L. Lovasz and J. Vygen, eds., *Research Trends in Combinatorial Optimization, Bonn 2008*, Springer-Verlag, Berlin, 2009, Chapter 11, 219–260.
- [9] H. Ohsugi and T. Hibi, Non-very ample configurations arising from contingency tables, preprint (2009). [arXiv:0904.3681v2](#). To appear in *Annals of the Institute of Statistical Mathematics*.
- [10] András Sebő, An introduction to empty lattice simplices. in: *Integer Programming and Combinatorial Optimization*, Lecture Notes in Computer Science **1610**, 400–414, Springer, Berlin, 1999.
- [11] G. M. Ziegler, *Lectures on Polytopes*, Springer-Verlag, New York, 1995.